

Hypothesis

Stage-variations of anandamide hydrolase activity in the mouse uterus during the natural oestrus cycle

Francesca G Klinger¹, Natalia Battista², Massimo De Felici*¹ and Mauro Maccarrone^{2,3}

Address: ¹Department of Public Health and Cell Biology, University of Rome Tor Vergata, 00133 Rome, Italy, ²Department of Biomedical Sciences, University of Teramo, 64100 Teramo, Italy and ³IRCCS C. Mondino, Mondino-Tor Vergata Center for Experimental Neurobiology, 00133 Rome, Italy

Email: Francesca G Klinger - klinger@uniroma2.it; Natalia Battista - battista@med.uniroma2.it; Massimo De Felici* - defelici@uniroma2.it; Mauro Maccarrone - Maccarrone@med.uniroma2.it

* Corresponding author

Published: 30 March 2006

Received: 28 November 2005

Accepted: 30 March 2006

Journal of Experimental & Clinical Assisted Reproduction 2006, **3**:3 doi:10.1186/1743-1050-3-3

This article is available from: <http://www.jexclinassistreprod.com/content/3/1/3>

© 2006 Klinger et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Recent studies have demonstrated that the endogenous cannabinoids are important modulators of fertility in mammals. In particular, a role of the endocannabinoid system in early stages of embryo development, oviductal transport of embryos, pregnancy maintenance and labour has been demonstrated in rodents and/or in humans. In the present paper, we report the analysis of FAAH activity and protein content in the mouse uterus as a function of the natural oestrus cycle stages. Variations of FAAH activity are discussed in relationship to changes in sex steroid levels and to the possible action of AEA on remodelling of uterine tissues.

Introduction

Recent studies have demonstrated that the endogenous cannabinoids (bioactive fatty acid amides and esters termed endocannabinoids) are important modulators of fertility in mammals [1,2]. The endocannabinoid anandamide (*N*-arachidonoylethanolamine, AEA) and congeners, cannabinoid CB1 and CB2 receptors, the AEA-synthesizing enzymes *N*-acetyltransferase (NAT) and *N*-acylphosphatidylethanolamine (NAPE)-specific phospholipase D, the AEA membrane transporter (AMT) and the AEA-hydrolyzing enzyme fatty acid amide hydrolase (FAAH), form the endocannabinoid system. A role of this system in early stages of embryo development [2], oviductal transport of embryos [3], pregnancy maintenance and labour [4,5], has been demonstrated in rodents and/or in humans.

During early pregnancy, a proper endocannabinoid signalling in the uterus and in the preimplantation embryo (both expressing CB1 receptor, AMT and FAAH), and possibly also in the circulating immune cells, appears to be crucial for embryo development and implantation [2]. In particular, high levels of AEA cause inhibition of trophoblast proliferation at the interimplantation sites and induce blastocyst apoptosis, while low levels of AEA at the implantation sites favour trophoblast differentiation and outgrowth [4,6-8]. In rodents, variation of AEA levels in the pregnant uterus has been directly correlated with FAAH expression and activity in uterine tissues [4,9,10]. Interestingly, FAAH activity and expression of at the mRNA level can be modulated in the pregnant uterus by the blastocyst itself, and by sex hormones [4,10,11]. Besides pregnant uterus and preimplantation embryo, oviduct is also a target for endocannabinoid action. In




Figure 1

Fluctuations of uterine FAAH in the normal oestrous cycle of mouse. FAAH activity and protein content were measured as reported in "Materials and Methods", and were expressed as percentage of the proestrus value (100% = 250 ± 30 pmol/min per mg protein, for the activity; 0.370 ± 0.030 absorbance units at 405 nm, for the protein content). Vertical bars represent S.D. values. Pro, proestrus; Est, oestrus; EMet, early metoestrus; LMet, late metoestrus; Di, dioestrus.

fact, in the mouse CB1 deficiency causes early pregnancy loss due to retention of embryos in the oviduct [3].

In the present paper, we report the analysis of FAAH activity and protein content in the mouse uterus as a function of the natural oestrous cycle stages. Variations of FAAH activity are discussed in relationship to changes in sex steroid levels and to the possible action of AEA on remodeling of uterine tissues.

Methods

Uterine tissue collection during the oestrous cycle

The 4–5 day oestrous cycle of mouse was monitored by the examination of vaginal smears and classified in five stages; proestrus, oestrus, early or late-metoestrus and dioestrus [12]. CD-1 mice were killed by cervical dislocation and uterus quickly removed in M2 without BSA [4], cut in small pieces and rapidly frozen and stored at -70°C.

Assay of FAAH activity and protein content

The hydrolysis of 10 µM [³H] AEA (223 Ci/mmol, from Perkin Elmer Life Sciences, Boston, MA) by the fatty acid amide hydrolase (E.C. 3.5.1.4; FAAH) activity was assayed in uterine extracts (20 µg/test)[4], by measuring the release of [³H] arachidonic acid, at pH 9.0, by reversed phase high performance liquid chromatography [4]. FAAH activity was expressed as pmol arachidonate released per min per mg protein.

FAAH protein content was determined in uterine homogenates (20 µg/test) by enzyme-linked immunosorbent assay (ELISA), performed as reported [13]. For the ELISA test rabbit anti-FAAH polyclonal antibodies [13] were prepared by Primm S.r.l. (Milan, Italy) and were used as first antibody (diluted 1:300), whereas goat anti-rabbit alkaline phosphatase conjugates (Bio-Rad Laboratories, Hercules, CA) were used as second antibodies (diluted 1:2000). Color development of the alkaline phosphatase reaction was measured at 405 nm, using *p*-nitrophenyl phosphate as substrate, and the ELISA test was linear in the range 0–50 µg/well of homogenate [13].

Data reported in this paper are the means ± S.D. of at least four independent experiments, each performed in duplicate. Statistical analysis was performed by the nonparametric Mann-Whitney U test, elaborating experimental data by means of the InStat 4 program (GraphPAD Software for Science, San Diego, CA).

Findings

The uterus undergoes cellular remodelling during each sexual cycle, in order to be ready for a possible pregnancy. In the absence of pregnancy, uterine changes are reversible permitting preparation in a subsequent cycle. In the event of mating and successful fertilization, however, the changes in the uterus take another route to support pregnancy. The uterine cellular changes during cycle and pregnancy are regulated by the circulating levels of ovarian sex steroids estradiol (E2) and progesterone (P).

The results of our analysis show that FAAH activity in the mouse uterus changed as function of the oestrous cycle stages, and that FAAH protein content paralleled the enzyme activity (Fig. 1). In particular, the highest FAAH activity and protein content were recorded at proestrus, followed by a progressive decrease at oestrus and metoestrus; finally, dioestrus was characterized by the lowest values for both parameters, down to ~20% or ~50% of the proestrus values respectively ($p < 0.01$ for dioestrus *versus* proestrus in both cases). In a variety of tissues high FAAH hydrolyzing activity has been associated with low AEA levels and *viceversa*, both in the central nervous system [14] and in the periphery [15]. In particular, in a previous study we have shown that uterine FAAH activity inversely correlates with uterine AEA levels, so that higher FAAH implicated lower AEA [11]. On this background, we can speculate that the levels of AEA are very likely to be low at proestrus, and that they progressively increase with a maximum at dioestrus.

Serum concentrations of estradiol (E2) and progesterone (P) in the CD-1 mouse oestrous cycle have been carefully analysed in previous studies [16,17]. It has been shown that there are two peaks in both E2 and P endogenous

concentrations. The highest E2 level occurs in proestrus, followed by a smaller peak in late metoestrus. The serum concentration of P exhibits a small peak in early oestrous and a large broad peak that spans both metoestrus and dioestrus. Following successful fertilization, both E2 and P are required for uterine receptivity and implantation.

By overlapping the literature data on the endogenous levels of E2 and P with the present analysis of FAAH, it appears that the highest levels of E2 and the lowest levels of P coincide with the highest FAAH activity (i.e., the lowest local levels of AEA); instead, a low FAAH activity (i.e., a high local level of AEA) parallels decreasing and increasing E2 and P concentrations, respectively. Altogether, these data suggest that in the non-pregnant uterus endogenous E2 and P exert opposite effects on FAAH activity, the first hormone being stimulatory and the second inhibitory. This seems at variance with the mouse pregnant uterus, where treatment with exogenous E2 and P down-modulates FAAH activity [4].

Among several effects that variation of FAAH activity and the resulting different local AEA levels could exert on the uterus, the action on the smooth muscle seems of particular interest. It has been demonstrated that endocannabinoids exert a potent and direct relaxant effect on human pregnant myometrium [18] and coordinate oviductal smooth muscle contraction during embryo transport [3]. Moreover, proliferation and apoptosis of uterine tissues are also possible targets of endocannabinoid action [19-22]. In fact, the pro-apoptotic activity of AEA has been documented in several cells and tissues [21], including mouse blastocyst [2]. It is well established that one of basic events of uterine cellular changes is the systemic cell turnover, that consists of cell renewal by proliferation and cell death by apoptosis. While uterine epithelial proliferative indices are highest on the proestrus day, epithelial apoptotic indices are lowest on the same day. A reverse pattern was demonstrated between uterine cell proliferation and apoptosis on the oestrous day [23-25]. The increased uterine cell proliferation has been correlated with increasing ovarian E2 secretion on the proestrus day [26], while increased uterine epithelial apoptosis on the oestrous day has been attributed to decline in serum E2 levels [27]. In addition, a number of studies in humans also showed that preovulatory ovarian estrogen secretion stimulates proliferation of uterine glandular epithelial cells, especially in functional endometrium, during proliferative phase. A rapid increase in the apoptotic index was noted in both epithelium and stroma during the last days of the cycle (late secretory phase), with a maximum index on the second day of menstruation [28]. On the basis of our study, it is possible to speculate that at proestrus low levels of AEA could favour the proliferation of uterine tis-

sues, while at metaestrus-dioestrus high levels of AEA contribute to programmed cell death.

In conclusion, we report here the analysis of FAAH activity and protein content in the mouse uterus, showing variations as a function of the oestrous cycle stages which can be related to changes of sex steroid levels and tissue remodelling. The present data may form the basis for further studies on the action of endocannabinoid signalling in this organ, and on its impact on reproductive physiology.

Competing interests

The author(s) declare that they have no competing interests.

Authors' contributions

FGK and NB have produced most of the results and have been involved in drafting of the manuscript. MDF and MM have made equal contributions to conception and design and analysis and interpretation of data as well. All authors have given final approval of the version of the manuscript to be published.

Acknowledgements

This work was supported by the Italian MIUR PRIN grant 2004-2005.

References

1. Park B, McPartland JM, Glass M: **Cannabis, cannabinoids and reproduction.** *Prostaglandins Leukot Essent Fatty Acids* 2004, **70**:189-97.
2. Maccarrone M, Finazzi-Agro A: **Anandamide hydrolase: a guardian angel of human reproduction?** *Trends Pharmacol Sci* 2004, **25**:353-7.
3. Wang H, Guo Y, Wang D, Kingsley PJ, Marnett LJ, Das SK, DuBois RN, Dey SK: **Aberrant cannabinoid signaling impairs oviductal transport of embryos.** *Nat Med* 2004, **10**:1074-80.
4. Maccarrone M, De Felici M, Bari M, Klinger F, Siracusa G, Finazzi-Agro A: **Down-regulation of anandamide hydrolase in mouse uterus by sex hormones.** *Eur J Biochem* 2000, **267**:2991-7.
5. Habayeb OM, Taylor AH, Evans MD, Cooke MS, Taylor DJ, Bell SC, Konje JC: **Plasma levels of the endocannabinoid anandamide in women – a potential role in pregnancy maintenance and labor?** *J Clin Endocrinol Metab* 2004, **89**:5482-7.
6. Paria BC, Deutsch DD, Dey SK: **The uterus is a potential site for anandamide synthesis and hydrolysis: differential profiles of anandamide synthase and hydrolase activities in the mouse uterus during the periimplantation period.** *Mol Reprod Dev* 1996, **45**:183-92.
7. Paria BC, Dey SK: **Ligand-receptor signaling with endocannabinoids in preimplantation embryo development and implantation.** *Chem Phys Lipids* 2000, **108**:211-20.
8. Liu WM, Duan EK, Cao YJ: **Effects of anandamide on embryo implantation in the mouse.** *Life Sci* 2002, **23**:1623-32.
9. Paria BC, Wang H, Dey SK: **Endocannabinoid signaling in synchronizing embryo development and uterine receptivity for implantation.** *Chem Phys Lipids* 2002, **121**:201-10.
10. Xiao AZ, Zhao YG, Duan EK: **Expression and regulation of the fatty acid amide hydrolase gene in the rat uterus during the estrous cycle and peri-implantation period.** *Mol Hum Reprod* 2002, **8**:651-8.
11. Maccarrone M, Frideri E, Bisogno T, Bari M, Cascio MG, Battista N, Finazzi Agro A, Suris R, Mechoulam R, Di Marzo V: **Up-regulation of the endocannabinoid system in the uterus of leptin knock-out (ob/ob) mice and implications for fertility.** *Mol Hum Reprod* 2005, **11**:21-8.

12. Bronson FH: **Male-induced precocial puberty in female mice: confirmation of the role of estrogen.** *Endocrinology* 1975, **96**:511-4.
13. Maccarrone M, Cecconi S, Rossi G, Battista N, Pauselli R, Finazzi-Agro A: **Anandamide activity and degradation are regulated by early postnatal aging and follicle-stimulating hormone in mouse Sertoli cells.** *Endocrinology* 2003, **144**:20-8.
14. Gubellini P, Picconi B, Bari M, Battista N, Calabresi P, Centonze D, Bernardi G, Finazzi-Agro A, Maccarrone M: **Experimental parkinsonism alters endocannabinoid degradation: implications for striatal glutamatergic transmission.** *J Neurosci* 2002, **22**:6900-7.
15. Lazzarin N, Valensise H, Bari M, Ubaldi F, Battista N, Finazzi-Agro A, Maccarrone M: **Fluctuations of fatty acid amide hydrolase and anandamide levels during the human ovulatory cycle.** *Gynecol Endocrinol* 2004, **18**:212-8.
16. Walmer DK, Wrona MA, Hughes CL, Nelson KG: **Lactoferrin expression in the mouse reproductive tract during the natural estrous cycle: correlation with circulating estradiol and progesterone.** *Endocrinology* 1992, **131**:1458-66.
17. Fata JE, Chaudhary V, Khokha R: **Cellular turnover in the mammary gland is correlated with systemic levels of progesterone and not 17beta-estradiol during the estrous cycle.** *Biol Reprod* 2001, **65**:680-8.
18. Denny MC, Friel AM, Houlihan DD, Broderick VM, Smith T, Morrison JJ: **Cannabinoids and the human uterus during pregnancy.** *Am J Obstet Gynecol* 2004, **190**:2-9.
19. Guzman M: **Cannabinoids: potential anticancer agents.** *Nat Rev Cancer* 2003, **3**:745-55.
20. Pomelli D: **The molecular logic of endocannabinoid signalling.** *Nat Rev Neurosci* 2003, **4**:873-84.
21. Maccarrone M, Finazzi-Agro A: **The endocannabinoid system, anandamide and the regulation of mammalian cell apoptosis.** *Cell Death Differ* 2003, **10**:946-55.
22. Di Marzo V, Bifulco M, De Petrocellis L: **The endocannabinoid system and its therapeutic exploitation.** *Nat Rev Drug Discov* 2004, **3**:771-84.
23. Huet-Hudson YM, Andrews GK, Dey SK: **Cell type-specific localization of c-myc protein in the mouse uterus: modulation by steroid hormones and analysis of the periimplantation period.** *Endocrinology* 1989, **125**:1683-90.
24. Pampfer S, Cordi S, Dutrieux C, Vanderheyden I, Marchand C, De Hertogh R: **Interleukin 1beta mediates the effect of high D-glucose on the secretion of TNF-alpha by mouse uterine epithelial cells.** *Cytokine* 1999, **11**:500-9.
25. Joswig A, Gabriel HD, Kibschull M, Winterhager E: **Apoptosis in uterine epithelium and decidua in response to implantation: evidence for two different pathways.** *Reprod Biol Endocrinol* 2003, **26**:1-44.
26. Short RV: **Role of hormones in sex cycle.** In *Reproduction In Mammals Volume 3.* Cambridge University Press; 1972:42-72.
27. Sato T, Nakajima H, Fujio K, Mori Y: **Enhancement of prostaglandin E2 production by epidermal growth factor requires the coordinate activation of cytosolic phospholipase A2 and cyclooxygenase 2 in human squamous carcinoma A431 cells.** *Prostaglandins* 1997, **53**:355-69.
28. Dahmoun M, Boman K, Cajander S, Westin P, Backstrom T: **Apoptosis, proliferation and sex hormone receptors in superficial parts of human endometrium at the end of the secretory phase.** *J Clin Endocrin Metab* 2005, **84**:1737-1743.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

